

PREVYMIS[®] (letermovir)

240 mg, 480 mg tablets

Injection 20 mg/mL

Oral pellets 20 mg, 120 mg per packet

If you're an adult who has received an allogeneic bone marrow transplant and are at high risk, **taking PREVYMIS can help prevent cytomegalovirus (CMV) infection and disease.**



Actor Portrayal

What is PREVYMIS?

PREVYMIS is a prescription medicine used to help prevent cytomegalovirus (CMV) infection and disease in adults who have received an allogeneic hematopoietic stem cell (bone marrow) transplant and who have a high risk for getting CMV infection and disease.

Important Safety Information About PREVYMIS

Do not take PREVYMIS if you take pimozide or ergot alkaloids.


If you are taking PREVYMIS with cyclosporine, do not take pitavastatin or simvastatin.

Important Safety Information continues on page 2.

Cytomegalovirus (CMV) is a common virus and almost anyone can get it

How do people get CMV?

CMV can spread through close contact with people who are infected with the virus. Infected people have the virus in their body fluids, such as saliva, urine, blood, breast milk, and semen. As people get older, they are more likely to have been in contact with CMV.



Over half of adults are infected with CMV by age 40.

What are the symptoms of CMV?

For most healthy adults, having CMV is not serious. Often there are no signs or symptoms of CMV. If there are symptoms, they can include fever, sore throat, and fatigue.

Does CMV go away?

Once a person gets CMV, it stays in the body for life. A healthy immune system can keep CMV from causing sickness. However, if the immune system is weakened, CMV can cause sickness and may also cause more serious symptoms.

If you've had a bone marrow transplant, you may have some questions about the impact of CMV

What does CMV have to do with a bone marrow transplant?

Some of the medicines that are needed for or after a bone marrow transplant can weaken the immune system. These include medicines that keep the body from rejecting a transplant. When your immune system is weakened, CMV may cause sickness.



Before a bone marrow transplant, people are tested for CMV. Laboratory tests, such as blood tests, can detect CMV in adults. With testing, healthcare providers know who has a greater risk for sickness from CMV.

CMV testing continues after a bone marrow transplant. The testing helps healthcare providers monitor CMV to determine if the virus is growing or multiplying.

How can CMV affect you after a bone marrow transplant?

CMV may become active after a bone marrow transplant and cause problems that can affect recovery. For example, sickness from CMV can affect the lungs, stomach, eyes, and liver.

CMV MANAGEMENT

Healthcare providers have ways to manage CMV. There are medicines that can help prevent CMV from becoming active.

Important Safety Information About PREVYMIS (continued)

Before taking PREVYMIS, tell your healthcare provider about all your medical conditions, including if you have kidney or liver problems; are pregnant or plan to become pregnant, since it is not known if PREVYMIS will harm your unborn baby; and if you are breastfeeding or plan to breastfeed, since it is not known if PREVYMIS passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking PREVYMIS.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. PREVYMIS may affect the way other medicines work, and other medicines may affect how PREVYMIS works and can cause serious side effects.

Important Safety Information continues on page 3.



About PREVYMIS

PREVYMIS is a prescription medicine used to help prevent cytomegalovirus (CMV) infection and disease in adults who:

- have received an allogeneic hematopoietic stem cell (bone marrow) transplant, **and**
- have a high risk of getting CMV infection and disease.



How does PREVYMIS work?

PREVYMIS helps prevent CMV from multiplying. That is how PREVYMIS helps keep people from getting sick from CMV.

How should I take PREVYMIS?

- PREVYMIS comes as a tablet, oral pellets, or can be given by your healthcare provider through an IV line (intravenously).
- PREVYMIS given intravenously should be used only if you are unable to take PREVYMIS tablets or PREVYMIS oral pellets. You should switch to PREVYMIS tablets or PREVYMIS oral pellets as soon as you are able to.
- Take PREVYMIS tablets or oral pellets exactly as your healthcare provider tells you to take it. Do not stop taking PREVYMIS without talking to your healthcare provider first.
- Take PREVYMIS tablets or oral pellets 1 time a day.
- It is important that you do not miss or skip doses of PREVYMIS.
- If you miss a dose of PREVYMIS tablets or oral pellets, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and take your dose at the next scheduled time. Do not take 2 doses of PREVYMIS at the same time or take more than your prescribed dose to make up for a missed dose.
- If you take too much PREVYMIS tablets or oral pellets, call your healthcare provider right away.



If you take PREVYMIS tablets:

- Take PREVYMIS with or without food.
- Swallow PREVYMIS tablets whole.



If you take PREVYMIS oral pellets:

- Your healthcare provider will tell you the number of PREVYMIS oral pellet packets to take for your prescribed dose.
- PREVYMIS oral pellets can be taken by mouth after mixing with soft food or given through a nasogastric tube (NG tube) or gastric tube (G tube).
- **See the Instructions for Use for the right way to prepare and take a dose of PREVYMIS oral pellets.** Keep the Instructions for Use and follow it each time you prepare and take PREVYMIS oral pellets.
- Do not crush or chew PREVYMIS oral pellets.



If you receive PREVYMIS through an IV line (intravenously):

- You will receive PREVYMIS 1 time each day given over 1 hour.
- If you miss or skip your dose of PREVYMIS, call your healthcare provider right away.

Important Safety Information About PREVYMIS (continued)

Know the medicines you take. Keep a list of medicines and show it to your healthcare provider and pharmacist when you get a new medicine. Your healthcare provider or pharmacist will tell you if it is safe to take PREVYMIS with other medicines. Do not start or stop taking another medicine without telling your healthcare provider first.

Important Safety Information continues on page 4.



Who should not take PREVYMIS?

Do not take PREVYMIS if you take:

- pimozide
- ergot alkaloids

If you are taking PREVYMIS with cyclosporine, do not take:

- pitavastatin or simvastatin

What should I tell my healthcare provider before taking PREVYMIS?

Before taking PREVYMIS, tell your healthcare provider about all your medical conditions, including if you:

- have kidney or liver problems.
- are pregnant or plan to become pregnant. It is not known if PREVYMIS will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if PREVYMIS passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking PREVYMIS.



Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements. PREVYMIS may affect the way other medicines work, and other medicines may affect how PREVYMIS works and cause serious side effects.

Know the medicines you take. Keep a list of medicines and show it to your healthcare provider and pharmacist when you get a new medicine. Your healthcare provider or pharmacist will tell you if it is safe to take PREVYMIS with other medicines. Do not start or stop taking another medicine without telling your healthcare provider first.



Actor Portrayal

Important Safety Information About PREVYMIS (continued)

The most common side effects of PREVYMIS in adults include: nausea, diarrhea, vomiting, swelling in your arms and legs, cough, headache, tiredness, and stomach (abdominal) pain.

These are not all the possible side effects of PREVYMIS.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Important Safety Information continues on page 5.

Talk to your healthcare provider and visit prevymis.com

to find out more about how PREVYMIS may help you.

Need more information about insurance coverage for PREVYMIS?



If you have been prescribed PREVYMIS, the Merck Access Program may be able to help:

- Identify your insurance benefits
- Obtain information about out-of-pocket costs
- Provide information on co-pay assistance options for eligible patients
- Refer you to the Merck Patient Assistance Program for eligibility determination (provided through the Merck Patient Assistance Program, Inc.)

Learn more at <https://www.merckaccessprogram-prevymis.com/hcc>

Questions?

Contact the Merck Access Program at 855-404-5278 Mon-Fri, 8 AM to 8 PM (ET), to speak to a representative.



Coupon offer for PREVYMIS

Eligible, privately insured patients may pay as little as \$15 per prescription on each of up to 8 qualifying prescriptions for up to a one-month's supply per prescription. Maximum savings is \$2,500 per prescription.^a

Click or scan the QR code to learn more.

^aNot valid for patients who are uninsured or patients with Medicare or other Government Program insurance. Not all patients are eligible. Certain restrictions apply. Visit prevymis.com for Terms and Conditions.

What is PREVYMIS?

PREVYMIS is a prescription medicine used to help prevent cytomegalovirus (CMV) infection and disease in adults who have received an allogeneic hematopoietic stem cell (bone marrow) transplant and who have a high risk for getting CMV infection and disease.

Important Safety Information About PREVYMIS

Do not take PREVYMIS if you take pimozone or ergot alkaloids.

If you are taking PREVYMIS with cyclosporine, do not take pitavastatin or simvastatin.

Please read the accompanying [Patient Information](#) and [Instructions for Use](#) for PREVYMIS and discuss it with your healthcare provider. The [Physician Prescribing Information](#) also is available.



Copyright © 2025 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. US-CYT-02160 03/25